

**Mpox can be treated by  
managing symptoms.  
Antiviral drugs are  
occasionally  
used for severe  
cases**



**If you think you  
have been exposed  
or have symptoms,  
please see a  
healthcare  
provider or call 811**



**apss**

### CONTACT US

AIDS PROGRAM SOUTH SASKATCHEWAN  
1325 Albert Street  
Regina, SK.  
306-924-8420  
[www.aidsprogramsouthsask.com](http://www.aidsprogramsouthsask.com)

Supported by:



**THIS MATERIAL WAS FUNDED BY A CONTRIBUTION FROM  
CBRC. THE VIEWS EXPRESSED HEREIN DO NOT  
NECESSARILY REPRESENT THE VIEWS OF CBRC.**

## **MPOX**


**Here's what you  
need to know**





## Transmission

- skin-to-skin contact
- touching bodily fluids of someone who has mpox
- contact with infected objects (ex. bedding, towels)
- large respiratory droplets




## Early Signs & Symptoms

It can take up to 21 days to develop symptoms.


- fever ( $>38^{\circ}\text{C}$ ), chills
- swollen lymph nodes
- headache
- muscle/joint pain
- backache
- sore throat
- cough
- stuffy nose
- exhaustion
- extreme weakness
- rash/blisters on face, palms of the hands, & soles of the feet within 1-3 days

## Prevention



Vaccine

Wear a mask & avoid close contact with infected individuals



Limit sex partners

Handwash



Disinfect objects & surfaces

